PRP for Neck Pain and Cervical Spine Conditions

23 SEP 2023 · 5 MIN READ

Neck pain can arise from discs, facets, nerves, ligaments, muscles and the shoulder. This guide explains why “PRP for neck pain” is not one procedure.

Key message: The cervical spine contains multiple possible pain generators beside critical neural and vascular structures. Treatment should be target-specific and based on high-quality imaging – not on the symptom “neck pain” alone.

What can cause neck pain?

Common structural sources include cervical facet joints, discs, nerve roots, uncovertebral joints, ligaments, muscles and post-traumatic change. Shoulder disease, headache disorders and peripheral nerve conditions may overlap. The same pain distribution can arise from different structures.

Can PRP be used in the cervical spine?

Cervical PRP must be discussed by target. For cervical facetogenic pain, a 2026 randomised double-blind trial found low-concentrate PRP and corticosteroid produced similar overall pain improvement across 6 months, although both groups improved and PRP showed some secondary benefits. For epidural PRP or platelet-derived products used for radicular pain, systematic reviews report potentially similar or longer-duration relief than steroid, but the certainty of evidence has been rated very low and much of the higher-level data are lumbar rather than cervical. [23,24]

Why advanced imaging matters

3T MRI can assess discs, cord, nerve roots, marrow and soft tissues. Dedicated nerve imaging may add information in selected neuropathic cases. CT can define bone, foraminal narrowing, calcification and complex post-surgical anatomy. Imaging should first rule out pathology that requires surgery or another specialist pathway.

Why precision is especially important in the neck

Cervical procedures occur close to the spinal cord, nerve roots and major blood vessels. The procedural approach and imaging modality must be chosen for the exact target. The potential benefit of any biologic must be weighed against target-specific procedural risk.

What should I expect after?

Local soreness or a temporary flare may occur. Some procedures can cause transient numbness or weakness if local anaesthetic is used nearby. Driving, work and activity instructions depend on the exact procedure and should be confirmed at booking.

When PRP is not the priority

Progressive weakness, spinal cord compression, unstable fracture, infection, tumour, severe myelopathy or a surgically remediable lesion should not be diverted into a generic regenerative pathway.

Frequently Asked Questions

Is PRP for neck pain the same as a cervical epidural?

No. A facet joint, epidural space, nerve root, ligament or tendon are different anatomical targets and have different evidence and risks.

Can MRI tell which neck structure hurts?

MRI identifies structural abnormalities and can narrow the possibilities, but clinical correlation is still required because many imaging findings can be asymptomatic.

Can PRP avoid neck surgery?

Not reliably. Some patients may improve with non-surgical treatment, but PRP should not delay surgery when neurological compression or instability makes surgery appropriate.

Important: New arm or leg weakness, gait disturbance, loss of hand coordination, bowel/bladder changes, fever, major trauma or severe rapidly progressive neurological symptoms require urgent assessment.

Related MBR Health resources

- PRP
- Pain & Injury
- 3T MRI
- Spectral CT
- Why Image-Guided PRP Matters
- PRP Aftercare

Related Patient Guides

Explore the complete MBR Health condition-specific PRP patient-guide series:

- PG8000 – PRP for Knee Osteoarthritis
- PG8001 – PRP for Hip Osteoarthritis
- PG8002 – PRP for Shoulder Tendon and Rotator Cuff Problems
- PG8003 – PRP for Tennis Elbow and Golfer's Elbow
- PG8004 – PRP for Achilles Tendinopathy
- PG8005 – PRP for Plantar Fasciitis
- PG8006 – PRP for Gluteal Tendinopathy and Greater Trochanteric Pain

- PG8007 – PRP for Hamstring, Adductor and Tendon Injuries
- PG8008 – PRP for Ligament Injuries
- PG8009 – PRP for Meniscus and Cartilage-Related Knee Pain
- PG8010 – PRP for Sacroiliac Joint Pain
- PG8011 – PRP for Facet Joint Pain
- PG8012 – PRP for Disc-Related Back Pain
- **PG8013 – PRP for Neck Pain and Cervical Spine Conditions (this guide)**
- PG8014 – PRP for Foot and Ankle Pain

For other FAQs, please visit the MBR Health FAQs page.

Looking for the Best Neck Pain Imaging and Precision Cervical Injection Clinic on the Gold Coast?

Patients comparing specialist neck-pain, cervical spine and regenerative injection clinics on the Gold Coast or Brisbane should ask whether the clinic can identify the actual pain generator before treatment. MBR Health combines advanced 3T MRI, Spectral CT, nerve-focused imaging and specialist radiologist-led image-guided procedures so cervical treatment is planned around anatomy, diagnosis and safety rather than a one-size-fits-all "neck injection".

References

[23] Allison DJ, Ebrahimzadeh S, Muise S, et al. Intra-articular corticosteroid injections versus platelet-rich plasma as a treatment for cervical facetogenic pain: a randomized clinical trial. Reg Anesth Pain Med. 2026;51(1):33-40. doi:10.1136/rapm-2024-105889.

[24] Kubrova E, Martinez Alvarez GA, Her YF, Pagan-Rosado R, Qu W, D'Souza RS. Platelet Rich Plasma and Platelet-Related Products in the Treatment of Radiculopathy-A Systematic Review of the Literature. Biomedicines. 2022;10(11):2813. doi:10.3390/biomedicines10112813.

[26] D'Souza RS, Her YF, Hussain N, et al. Evidence-Based Clinical Practice Guidelines on Regenerative Medicine Treatment for Chronic Pain: A Consensus Report from a Multispecialty Working Group. J Pain Res. 2024;17:2951-3001. doi:10.2147/JPR.S480559.

General information only. PRP suitability, medication changes, dosing, procedure choice and rehabilitation require individual clinical assessment. Evidence and regulatory requirements continue to evolve.

Read this article online: https://mbrhealth.com.au/insights/prp-for-neck-pain-and-cervical-spine-conditions/

© 2026 MBR Health · mbrhealth.com.au